



**ISPM 28**

**INTERNATIONAL STANDARDS FOR  
PHYTOSANITARY MEASURES**

**ISPM 28**

**PHYTOSANITARY TREATMENTS FOR  
REGULATED PESTS**

**(2007)**

Produced by the Secretariat of the International Plant Protection Convention



**Publication history**

*This is not an official part of the standard*

2004-04 ICPM-6 added topic *Phytosanitary treatments for regulated pests* (2004-028)

2004-11 SC approved Specification 22 *Research protocols for phytosanitary measures*

2005-08 TPPT developed draft text and sent for MC

2005-10 MC under fast-track process

2005-11 SC requested further review

2006-05 SC revised draft text and approved for MC

2006-06 Sent for MC

2006-11 SC revised draft text

2007-03 CPM-2 adopted standard

**ISPM 28.** 2007. *Phytosanitary treatments for regulated pests*. Rome, IPPC, FAO.

Publication histories of attachments are included in each attachment

Publication history: Last modified August 2011

**CONTENTS**

Adoption.....	28-5
INTRODUCTION.....	28-5
Scope.....	28-5
References.....	28-5
Definitions.....	28-5
Outline of Requirements.....	28-5
BACKGROUND.....	28-7
REQUIREMENTS.....	28-7
1. Purpose and Use.....	28-7
2. Process for Treatment Submission and Adoption.....	28-8
3. Requirements for Phytosanitary Treatments.....	28-8
3.1 Summary information.....	28-9
3.2 Efficacy data in support of the submission of a phytosanitary treatment.....	28-9
3.2.1 Efficacy data under laboratory/controlled conditions.....	28-9
3.2.2 Efficacy data using operational conditions.....	28-10
3.3 Feasibility and applicability.....	28-11
4. Evaluation of Submitted Treatments.....	28-11
5. Publication of Phytosanitary Treatments.....	28-12
6. Treatment Review and Re-evaluations.....	28-12
APPENDIX 1 (2011): Lists of adopted annexes.....	28-13



## Adoption

This standard was adopted by the Second Session of the Commission on Phytosanitary Measures in March 2007. Adoption information for attachments is stated in each attachment, if different from core text.

## INTRODUCTION

### Scope

This standard presents as annexes phytosanitary treatments evaluated and adopted by the Commission on Phytosanitary Measures (CPM). It also describes the requirements for submission and evaluation of the efficacy data and other relevant information on a phytosanitary treatment that can be used as a phytosanitary measure and that will be annexed to this standard after its adoption.

The treatments are for the control of regulated pests on regulated articles, primarily those moving in international trade. The adopted treatments provide the minimum requirements necessary to control a regulated pest at a stated efficacy.

The scope of this standard does not include issues related to pesticide registration or other domestic requirements for approval of treatments (e.g. irradiation)<sup>1</sup>.

### References

**IPPC.** 1997. *International Plant Protection Convention*. Rome, IPPC, FAO.

**ISPM 5.** *Glossary of phytosanitary terms*. Rome, IPPC, FAO.

**ISPM 11.** 2004. *Pest risk analysis for quarantine pests including analysis of environmental risks and living modified organisms*. Rome, IPPC, FAO.

### Definitions

Definitions of phytosanitary terms used in the present standard can be found in ISPM 5 (*Glossary of phytosanitary terms*).

### Outline of Requirements

Harmonized phytosanitary treatments support efficient phytosanitary measures in a wide range of circumstances and enhance the mutual recognition of treatment efficacy. Annexes to this standard contain those phytosanitary treatments which have been adopted by the CPM.

National plant protection organizations (NPPOs) and regional plant protection organizations (RPPOs) may submit data and other information for the evaluation of efficacy, feasibility and applicability of treatments. The information should include a detailed description of the treatment, including efficacy data, the name of a contact person and the reason for the submission. Treatments that are eligible for evaluation include mechanical, chemical, irradiation, physical and controlled atmosphere treatments. The efficacy data should be clear and should preferably include data on the treatment under laboratory or controlled conditions as well as under operational conditions. Information on feasibility and applicability of the proposed treatment(s) should include items on cost, commercial relevance, level of expertise required to apply the treatment and versatility.

---

<sup>1</sup> The inclusion of a phytosanitary treatment in this ISPM does not create any obligation for a contracting party to approve the treatment or register or adopt it for use in its territory.

Submissions with complete information will be considered by the Technical Panel on Phytosanitary Treatments (TPPT), and if the treatment is deemed acceptable, it will be recommended to the CPM for adoption.

## BACKGROUND

The purpose of the IPPC is “to prevent the spread and introduction of pests of plants and plant products, and to promote appropriate measures for their control” (Article I.1 of the IPPC). The requirement or application of phytosanitary treatments to regulated articles is a phytosanitary measure used by contracting parties to prevent the introduction and spread of regulated pests.

Article VII.1 of the IPPC 1997 states:

contracting parties shall have sovereign authority to regulate, in accordance with applicable international agreements, the entry of plants and plant products and other regulated articles and, to this end, may:

- (a) prescribe and adopt phytosanitary measures concerning the importation of plants, plant products and other regulated articles, including, for example, inspection, prohibition on importation, and treatment.

Phytosanitary measures required by a contracting party shall be technically justified (Article VII.2(a) of the IPPC).

Phytosanitary treatments are used by NPPOs to prevent the introduction and spread of regulated pests. Many of these treatments are supported by extensive research data, and others are used based on historical evidence supporting their efficacy. In practice, many countries use the same treatments or similar treatments for specified pests; however, mutual recognition is often a complex and difficult process. Furthermore, there has previously been neither an internationally recognized organization or process to evaluate treatments for their efficacy nor a central repository for listing such treatments. The Interim Commission on Phytosanitary Measures, at its sixth session in 2004, recognized the need for international recognition of phytosanitary treatments of major importance and approved the formation of the TPPT for that purpose.

## REQUIREMENTS

### 1. Purpose and Use

The purpose of harmonizing phytosanitary treatments is to support efficient phytosanitary measures in a wide range of circumstances and to enhance the mutual recognition of treatment efficacy by NPPOs, which may also facilitate trade. Furthermore, these treatment schedules should aid the development of expertise and technical cooperation. NPPOs are not obliged to use these treatments and may use other phytosanitary treatments for treating the same regulated pests or regulated articles.

Adopted phytosanitary treatments provide a means for the killing, inactivation or removal of pests, for rendering pests infertile or for devitalization, at a stated efficacy, and are relevant primarily to international trade. The level of efficacy, specificity and applicability of each treatment is indicated where possible. NPPOs may use these criteria to select the treatment or combination of treatments that are appropriate for the relevant circumstances.

When requiring phytosanitary treatments for imports, contracting parties should take into account the following points:

- Phytosanitary measures required by a contracting party shall be technically justified.
- Phytosanitary treatments contained in annexes of this standard have the status of an ISPM and therefore should be considered accordingly.
- Regulatory regimes of exporting contracting parties may prevent certain treatments from being approved for use within their territories. Therefore efforts should be made to accept equivalent treatments where possible.

## 2. Process for Treatment Submission and Adoption

The submission process is initiated by a call for topics for standards (including topics for treatments) according to the “IPPC standard setting procedure” and the “Procedure and criteria for identifying topics for inclusion in the IPPC standard setting work programme”. These procedures are provided on the International Phytosanitary Portal (<https://www.ippc.int>).

In particular, the following points apply to treatments:

- Once a topic for treatments (e.g. treatments for fruit flies or for pests on wood) has been added to the IPPC standard setting work programme, the IPPC Secretariat, under direction of the Standards Committee (with recommendations from the TPPT), will call for the submissions and data on treatments on that topic.
- NPPOs or RPPOs submit treatments (accompanied by relevant information as requested in section 3) to the Secretariat.
- Only submissions of treatments that are deemed by the NPPO or RPPO to meet the requirements listed in this standard should be submitted, and it is recommended that these treatments have been approved for national use before their submission. Treatments include, but are not limited to, mechanical, chemical, irradiation, physical (heat, cold) and controlled atmosphere treatments. NPPOs and RPPOs should take into account other factors when considering phytosanitary treatments for submission, such as the effects on human health and safety, animal health and the impact on the environment (as described in the preamble and Article I.1 of the IPPC and in Article III of the IPPC regarding relationship with other international agreements). Effects on the quality and intended use of the regulated article should also be considered.
- Treatment submissions will be evaluated based on the requirements listed in section 3. If large numbers of submissions are received, the TPPT will work with the Standards Committee to determine the priority for reviewing submissions.
- Treatments that meet the requirements listed in section 3 will be recommended and the treatment submitted, along with a report and a summary of the information evaluated, to the Standards Committee and in turn to the IPPC standard setting process. The report of the technical panel with the summary information and the SC report will be available to contracting parties. Further detailed information (as long as it is not confidential) will be available on request from the Secretariat.
- The CPM will adopt or reject a treatment. If adopted, the treatment is annexed to this standard.

## 3. Requirements for Phytosanitary Treatments

For the purpose of this standard, phytosanitary treatments should fulfil the following requirements:

- be effective in killing, inactivating or removing pests, or rendering pests infertile or for devitalization associated with a regulated article. The level of efficacy of the treatment should be stated (quantified or expressed statistically). Where experimental data is unavailable or insufficient, other evidence that supports the efficacy (i.e. historical and/or practical information/experience) should be provided.
- be well documented to show that the efficacy data has been generated using appropriate scientific procedures, including where relevant an appropriate experimental design. The data supporting the treatment should be verifiable, reproducible, and based on statistical methods and/or on established and accepted international practice; preferably the research should have been published in a peer-reviewed journal.
- be feasible and applicable for use primarily in international trade or for other purposes (e.g. to protect endangered areas domestically, or for research).
- not be phytotoxic or have other adverse effects.

Submissions of phytosanitary treatments should include the following:

- summary information
- efficacy data in support of the phytosanitary treatment
- information on feasibility and applicability.

### **3.1 Summary information**

The summary information should be submitted by NPPOs or RPPOs to the Secretariat and should include:

- name of the treatment
- name of the NPPO or RPPO and contact information
- name and contact details of a person responsible for submission of the treatment
- treatment description (active ingredient(s), treatment type, target regulated article(s), target pest(s), treatment schedule, and other relevant information)
- reason for submission, including its relevance to existing ISPMs.

Submissions should utilize a form provided by the IPPC Secretariat and available on the International Phytosanitary Portal (<https://www.ippc.int>).

In addition, the NPPO or RPPO should describe the experience or expertise in the subject area of the laboratory, organization and/or scientist(s) involved in producing the data, and any quality assurance system or accreditation programme applied in the development and/or testing of the phytosanitary treatment. This information will be considered when evaluating the data submitted.

### **3.2 Efficacy data in support of the submission of a phytosanitary treatment**

The source of all efficacy data (published or unpublished) should be provided in the submission. Supporting data should be presented clearly and systematically. Any claims on the efficacy must be substantiated by data.

#### **3.2.1 Efficacy data under laboratory/controlled conditions**

The life-cycle stage of the target pest for the treatment should be specified. Usually, the life stage(s) associated with the regulated article moving in trade is the stage for which a treatment is proposed and established. In some circumstances, e.g. where several life stages may occur on the regulated article, the most resistant life stage of the pest should be used for testing a treatment. However, practical considerations should be taken into account, as well as pest control strategies aimed at exploiting more vulnerable or otherwise specific stages of a pest. If efficacy data is submitted for a life stage that is not considered to be the most resistant (e.g. if the most resistant life stage is not associated with the regulated article), rationale for this should be provided. The efficacy data provided should specify the statistical level of confidence supporting efficacy claims made for treatment of the specified life stage.

Where possible, data should be presented on methods used to determine the effective dose/treatment to demonstrate the range of efficacy of the treatment (e.g. dose/efficacy curves). Treatments can normally be evaluated only for the conditions under which they were tested. However, additional information can be provided to support any extrapolation if the scope of a treatment is to be extended (e.g. extension of the range of temperatures, inclusion of other cultivars or pest species). Where the information provided is adequate to demonstrate the effectiveness of the treatment, only a summary of relevant preliminary laboratory tests will be required. The materials and methods used in the experiments should be suitable for the use of the treatment at the stated efficacy.

The data provided should include detailed information on, but not limited to, the following elements:

#### **Pest information**

- identity of the pest to the appropriate level (e.g. genus, species, strain, biotype, physiological race), life stage, and if laboratory or field strain was used
- conditions under which the pests are cultured, reared or grown
- biological traits of the pest relevant to the treatment (e.g. viability, genetic variability, weight, developmental time, development stage, fecundity, freedom from disease or parasites)
- method of natural or artificial infestation
- determination of most resistant species/life stage (in the regulated article where appropriate)

#### **Regulated article information**

- type of regulated article and intended use
- botanical name for plant or plant product (where applicable)
  - . type/cultivar. A requirement for varietal testing should be based on evidence that the varietal differences impact treatment efficacy, and data should be provided to support the requirement.
- conditions of the plant or plant product, for example:
  - . whether it was free from non-target pest infestation, non-pest disorder or pesticide residue
  - . size, shape, weight, stage of maturity, quality etc.
  - . whether infested at a susceptible growth stage
  - . storage conditions after harvest

#### **Experimental parameters**

- level of confidence of laboratory tests provided by the method of statistical analysis and the data supporting that calculation (e.g. number of subjects treated, number of replicate tests, controls)
- experimental facilities and equipment
- experimental design (e.g. randomized complete block design) if needed
- experimental conditions (e.g. temperature, relative humidity, diurnal cycle)
- monitoring of critical parameters (e.g. exposure time, dose, temperature of regulated article and ambient air, relative humidity)
- methodology to measure the effectiveness of the treatment (e.g. whether mortality is the proper parameter, whether the end-point mortality was assessed at the correct time, the mortality or sterility of the treated and control groups)
- determination of efficacy over a range of critical parameters, where appropriate, such as exposure time, dose, temperature, relative humidity and water content, size and density
- methodology to measure phytotoxicity, when appropriate
- dosimetry system, calibration and accuracy of measurements, if using irradiation.

### **3.2.2 Efficacy data using operational conditions**

Treatments may be submitted for evaluation without going through the processes outlined in section 3.2.1 when there is sufficient efficacy data available from the operational application of the treatment. When a treatment has been developed under laboratory conditions, it should be validated by testing under operational or simulated operational conditions. Results of these tests should confirm that the application of the treatment schedule achieves the stated efficacy under conditions in which the treatment will be used.

Where treatment specifications differ for trials under operational conditions, the test protocol modifications should be indicated. Supporting data may be presented from preliminary tests to refine

the treatment schedule to establish the effective dose (e.g. temperature, chemical, irradiation) under operational conditions.

In some cases the method of achieving the effective dose will be different from the method established under laboratory conditions. Data that supports any extrapolation of laboratory results should be provided.

The same data requirements as listed in section 3.2.1 should also be provided for these tests. Other data required, depending on whether the treatments are carried out pre- or post-harvest, are listed below:

- factors that affect the efficacy of the treatment (e.g. for post-harvest treatments: packaging, packing method, stacking, timing of treatments (pre/post packaging or processing, in transit, on arrival)). The circumstances of the treatment should be stated, for example the efficacy of a treatment may be affected by packaging, and data should be provided to support all the circumstances that are applicable.
- monitoring of critical parameters (e.g. exposure time, dose, temperature of regulated article and ambient air, relative humidity). For example:
  - . the number and placement of gas sampling lines (fumigation)
  - . the number and placement of temperature/humidity sensors.

In addition, any special procedures that affect the success of the treatment (e.g. to maintain the quality of the regulated article) should be included.

### **3.3 Feasibility and applicability**

Information should be provided, where appropriate, to evaluate if the phytosanitary treatment is feasible and applicable. This includes such items as:

- procedure for carrying out the phytosanitary treatment (including ease of use, risks to operators, technical complexity, training required, equipment required, facilities needed)
- cost of typical treatment facility and operational running costs if appropriate
- commercial relevance, including affordability
- extent to which other NPPOs have approved the treatment as a phytosanitary measure
- availability of expertise needed to apply the phytosanitary treatment
- versatility of the phytosanitary treatment (e.g. application to a wide range of countries, pests and commodities)
- the degree to which the phytosanitary treatment complements other phytosanitary measures (e.g. potential for the treatment to be used as part of a systems approach for one pest or to complement treatments for other pests)
- summary of available information of potential undesirable side-effects (e.g. impacts on the environment, impacts on non-target organisms, human and animal health)
- applicability of treatment with respect to specific regulated article/pest combinations
- technical viability
- phytotoxicity and other effects on the quality of regulated articles, when appropriate
- consideration of the risk of the target organism having or developing resistance to the treatment.

Treatment procedures should adequately describe the method for applying the treatment in a commercial setting.

## **4. Evaluation of Submitted Treatments**

Submissions will be considered by the TPPT only when the information outlined in section 3 is fully addressed. The information provided will be evaluated against the requirements in section 3.

Due respect for confidentiality will be exercised when the confidential nature of information is indicated. In such cases, the confidential information within the submission should be clearly identified. Where confidential information is essential for the adoption of the treatment, the submitter will be requested to release the information. If the release of the information is not granted, the adoption of the treatment may be affected.

Treatments will be adopted only for the regulated articles and target species for which they were tested and for the conditions under which they were tested, unless data is presented to support extrapolation (e.g. to apply the treatment to a range of pest species or regulated articles).

If the submission fails to meet the requirements outlined in section 3, the reason(s) will be communicated to the contact identified on the submission. There may be a recommendation to provide additional information or to initiate further work (e.g. research, field testing, analysis).

## **5. Publication of Phytosanitary Treatments**

After adoption by the CPM, phytosanitary treatments will be annexed to this standard. (Appendix 1 provides lists of the adopted annexes.)

## **6. Treatment Review and Re-evaluations**

Contracting parties should submit to the IPPC Secretariat any new information that could have an impact on the treatments currently adopted by the CPM. The TPPT will review the data and revise the treatments if necessary through the normal standard setting process.

Appendix 1 is for reference purposes only and is not an official part of the standard.

This appendix was updated by the Secretariat in September 2011.

## APPENDIX 1: Lists of adopted annexes

The following phytopsanitary treatments for regulated pests have been adopted by the Commission of Phytopsanitary Measures as annexes to ISPM 28:2007. These phytopsanitary treatments are available as separate documents on the International Phytopsanitary Portal (<https://www.ippc.int>).

Annexes are listed by organism, by regulated article and by treatment type.

### Adopted annexes by target pest

Target pest	Taxonomic information	Target regulated articles	Treatment type	Treatment schedule (e.g. active ingredient, dose)	Annex no. (PT no.)	Adoption year
<i>Anastrepha ludens</i>	Diptera: Tephritidae	Fruits and vegetables	Irradiation	70 Gy (Minimum absorbed dose)	1	2009
<i>Anastrepha obliqua</i>	Diptera: Tephritidae	Fruits and vegetables	Irradiation	70 Gy (Minimum absorbed dose)	2	2009
<i>Anastrepha serpentina</i>	Diptera: Tephritidae	Fruits and vegetables	Irradiation	100 Gy (Minimum absorbed dose)	3	2009
<i>Bactrocera jarvisi</i>	Diptera: Tephritidae	Fruits and vegetables	Irradiation	100 Gy (Minimum absorbed dose)	4	2009
<i>Bactrocera tryoni</i>	Diptera: Tephritidae	Fruits and vegetables	Irradiation	100 Gy (Minimum absorbed dose)	5	2009
<i>Ceratitis capitata</i>	Diptera: Tephritidae	Fruits and vegetables	Irradiation	100 Gy (Minimum absorbed dose)	14	2011
<i>Conotrachelus nenuphar</i>	Coleoptera: Curculionidae	Fruits and vegetables	Irradiation	92 Gy (Minimum absorbed dose)	9	2010
<i>Cydia pomonella</i>	Lepidoptera: Tortricidae	Fruits and vegetables	Irradiation	200 Gy (Minimum absorbed dose)	6	2009
<i>Cylas formicarius elegantulus</i>	Coleoptera: Brentidae	Fruits and vegetables	Irradiation	165 Gy (Minimum absorbed dose)	12	2011
<i>Euscepes postfasciatus</i>	Coleoptera: Curculionidae	Fruits and vegetables	Irradiation	150 Gy (Minimum absorbed dose)	13	2011
Fruit flies of the family Tephritidae (generic)	Diptera: Tephritidae	Fruits and vegetables	Irradiation	150 Gy (Minimum absorbed dose)	7	2009
<i>Grapholita molesta</i>	Lepidoptera: Tortricidae	Fruits and vegetables	Irradiation	232 Gy (Minimum absorbed dose)	10	2010
<i>Grapholita molesta</i> under hypoxia	Lepidoptera: Tortricidae	Fruits and vegetables	Irradiation	232 Gy (Minimum absorbed dose)	11	2010
<i>Rhagoletis pomonella</i>	Diptera: Tephritidae	Fruits and vegetables	Irradiation	60 Gy (Minimum absorbed dose)	8	2009

**Adopted annexes by target regulated articles**

Target regulated articles	Target pest	Taxonomic information	Treatment type	Treatment schedule (e.g. active ingredient, dose)	Annex no. (PT no.)	Adoption year
Fruits and vegetables	<i>Anastrepha ludens</i>	Diptera: Tephritidae	Irradiation	70 Gy (Minimum absorbed dose)	1	2009
Fruits and vegetables	<i>Anastrepha obliqua</i>	Diptera: Tephritidae	Irradiation	70 Gy (Minimum absorbed dose)	2	2009
Fruits and vegetables	<i>Anastrepha serpentina</i>	Diptera: Tephritidae	Irradiation	100 Gy (Minimum absorbed dose)	3	2009
Fruits and vegetables	<i>Bactrocera jarvisi</i>	Diptera: Tephritidae	Irradiation	100 Gy (Minimum absorbed dose)	4	2009
Fruits and vegetables	<i>Bactrocera tryoni</i>	Diptera: Tephritidae	Irradiation	100 Gy (Minimum absorbed dose)	5	2009
Fruits and vegetables	<i>Ceratitis capitata</i>	Diptera: Tephritidae	Irradiation	100 Gy (Minimum absorbed dose)	14	2011
Fruits and vegetables	<i>Conotrachelus nenuphar</i>	Coleoptera: Curculionidae	Irradiation	92 Gy (Minimum absorbed dose)	9	2010
Fruits and vegetables	<i>Cydia pomonella</i>	Lepidoptera: Tortricidae	Irradiation	200 Gy (Minimum absorbed dose)	6	2009
Fruits and vegetables	<i>Cylas formicarius elegantulus</i>	Coleoptera: Brentidae	Irradiation	165 Gy (Minimum absorbed dose)	12	2011
Fruits and vegetables	<i>Euscepes postfasciatus</i>	Coleoptera: Curculionidae	Irradiation	150 Gy (Minimum absorbed dose)	13	2011
Fruits and vegetables	Fruit flies of the family Tephritidae (generic)	Diptera: Tephritidae	Irradiation	150 Gy (Minimum absorbed dose)	7	2009
Fruits and vegetables	<i>Grapholita molesta</i>	Lepidoptera: Tortricidae	Irradiation	232 Gy (Minimum absorbed dose)	10	2010
Fruits and vegetables	<i>Grapholita molesta</i> under hypoxia	Lepidoptera: Tortricidae	Irradiation	232 Gy (Minimum absorbed dose)	11	2010
Fruits and vegetables	<i>Rhagoletis pomonella</i>	Diptera: Tephritidae	Irradiation	60 Gy (Minimum absorbed dose)	8	2009

**Adopted annexes by treatment type**

Treatment type	Target pest	Taxonomic information	Target regulated articles	Treatment schedule (e.g. active ingredient, dose)	Annex no. (PT no.)	Adoption year
Irradiation	<i>Anastrepha ludens</i>	Diptera: Tephritidae	Fruits and vegetables	70 Gy (Minimum absorbed dose)	1	2009
Irradiation	<i>Anastrepha obliqua</i>	Diptera: Tephritidae	Fruits and vegetables	70 Gy (Minimum absorbed dose)	2	2009
Irradiation	<i>Anastrepha serpentina</i>	Diptera: Tephritidae	Fruits and vegetables	100 Gy (Minimum absorbed dose)	3	2009
Irradiation	<i>Bactrocera jarvisi</i>	Diptera: Tephritidae	Fruits and vegetables	100 Gy (Minimum absorbed dose)	4	2009
Irradiation	<i>Bactrocera tryoni</i>	Diptera: Tephritidae	Fruits and vegetables	100 Gy (Minimum absorbed dose)	5	2009

Irradiation	<i>Ceratitis capitata</i>	Diptera: Tephritidae	Fruits and vegetables	100 Gy (Minimum absorbed dose)	<b>14</b>	2011
Irradiation	<i>Conotrachelus nenuphar</i>	Coleoptera: Curculionidae	Fruits and vegetables	92 Gy (Minimum absorbed dose)	<b>9</b>	2010
Irradiation	<i>Cydia pomonella</i>	Lepidoptera: Tortricidae	Fruits and vegetables	200 Gy (Minimum absorbed dose)	<b>6</b>	2009
Irradiation	<i>Cylas formicarius elegantulus</i>	Coleoptera: Brentidae	Fruits and vegetables	165 Gy (Minimum absorbed dose)	<b>12</b>	2011
Irradiation	<i>Euscepes postfasciatus</i>	Coleoptera: Curculionidae	Fruits and vegetables	150 Gy (Minimum absorbed dose)	<b>13</b>	2011
Irradiation	Fruit flies of the family Tephritidae (generic)	Diptera: Tephritidae	Fruits and vegetables	150 Gy (Minimum absorbed dose)	<b>7</b>	2009
Irradiation	<i>Grapholita molesta</i>	Lepidoptera: Tortricidae	Fruits and vegetables	232 Gy (Minimum absorbed dose)	<b>10</b>	2010
Irradiation	<i>Grapholita molesta</i> under hypoxia	Lepidoptera: Tortricidae	Fruits and vegetables	232 Gy (Minimum absorbed dose)	<b>11</b>	2010
Irradiation	<i>Rhagoletis pomonella</i>	Diptera: Tephritidae	Fruits and vegetables	60 Gy (Minimum absorbed dose)	<b>8</b>	2009



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 1: Irradiation treatment for *Anastrepha ludens* (2009)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 70 Gy minimum absorbed dose to prevent the emergence of adults of *Anastrepha ludens* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Anastrepha ludens</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Anastrepha ludens</i> (Loew) (Diptera: Tephritidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Anastrepha ludens</i>

##### **Treatment schedule**

Minimum absorbed dose of 70 Gy to prevent the emergence of adults of *Anastrepha ludens*.

Efficacy and confidence level of the treatment is ED<sub>99,9968</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18 (*Guidelines for the use of irradiation as a phytosanitary measure*).

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Anastrepha ludens* (larvae and/or pupae) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Hallman & Martinez (2001) that determined the efficacy of irradiation as a treatment for this pest in *Citrus paradisi*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J. 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L. 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J. 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R. 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M. 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M. 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L. 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A. 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

**Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for Anastrepha ludens* (2006--130)

2006-12 TPPT developed draft text

2007-05 SC approved draft text for MC

2007-10 Sent for MC under fast-track process

2008-07 TPPT revised draft text

2008-12 SC revised draft text for adoption via e-decision

2009-03 CPM-4 adopted Annex 1 to ISPM 28

**ISPM 28. 2007: Annex 1** *Irradiation treatment for Anastrepha ludens* (2009).  
Rome, IPPC, FAO.

Publication history: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 2: Irradiation treatment for *Anastrepha obliqua* (2009)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 70 Gy minimum absorbed dose to prevent the emergence of adults of *Anastrepha obliqua* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Anastrepha obliqua</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Anastrepha obliqua</i> (Macquart) (Diptera: Tephritidae)
<b>Target regulated articles:</b>	All fruits and vegetables, including nuts that are hosts of <i>Anastrepha obliqua</i> .

##### **Treatment schedule**

Minimum absorbed dose of 70 Gy to prevent the emergence of adults of *Anastrepha obliqua*.

Efficacy and confidence level of the treatment is ED<sub>99,9968</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18 (*Guidelines for the use of irradiation as a phytosanitary measure*).

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Anastrepha obliqua* (larvae and/or pupae) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Bustos *et al.* (2004), Hallman & Martinez (2001) and Hallman & Worley (1999) that determined the efficacy of irradiation as a treatment for this pest in *Citrus paradisi* and *Mangifera indica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Hallman, G.J. & Worley, J.W.** 1999. Gamma radiation doses to prevent adult emergence from immatures of Mexican and West Indian fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 92: 967–973.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

**Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for Anastrepha obliqua* (2006-115)

2006-12 TPPT developed draft text

2007-05 SC approved draft text for MC

2007-10 Sent for MC under fast-track process

2008-07 TPPT revised draft text

2008-12 SC revised draft for adoption via e-decision

2009-03 CPM-4 adopted Annex 2 to ISPM 28

**ISPM 28.** 2007: **Annex 2** *Irradiation treatment for Anastrepha obliqua* (2009).  
Rome, IPPC, FAO.

Publication history: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 3: Irradiation treatment for *Anastrepha serpentina* (2009)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 100 Gy minimum absorbed dose to prevent the emergence of adults of *Anastrepha serpentina* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Anastrepha serpentina</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Anastrepha serpentina</i> (Wiedemann) (Diptera: Tephritidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Anastrepha serpentina</i> .

##### **Treatment schedule**

Minimum absorbed dose of 100 Gy to prevent the emergence of adults of *Anastrepha serpentina*.

Efficacy and confidence level of the treatment is ED<sub>99,9972</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18 (*Guidelines for the use of irradiation as a phytosanitary measure*).

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Anastrepha serpentina* (larvae and/or pupae) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Bustos *et al.* (2004) that determined the efficacy of irradiation as a treatment for this pest in *Mangifera indica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J. 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L. 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J. 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R. 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M. 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M. 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L. 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A. 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

**Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for Anastrepha serpentina* (2006-116)

2006-12 TPPT developed draft text

2007-05 SC approved draft text for MC

2007-10 Sent for MC under fast-track process

2008-07 TPPT revised draft text

2008-12 SC revised draft text for adoption via e-decision

2009-03 CPM-4 adopted Annex 3 to ISPM 28

**ISPM 28. 2007: Annex 3** *Irradiation treatment for Anastrepha serpentina* (2009).  
Rome, IPPC, FAO.

Publication notes: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 4: Irradiation treatment for *Bactrocera jarvisi* (2009)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 100 Gy minimum absorbed dose to prevent the emergence of adults of *Bactrocera jarvisi* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Bactrocera jarvisi</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Bactrocera jarvisi</i> (Tryon) (Diptera: Tephritidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Bactrocera jarvisi</i> .

##### **Treatment schedule**

Minimum absorbed dose of 100 Gy to prevent the emergence of adults of *Bactrocera jarvisi*.

Efficacy and confidence level of the treatment is ED<sub>99,9981</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18 (*Guidelines for the use of irradiation as a phytosanitary measure*).

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Bactrocera jarvisi* (larvae and/or pupae) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Heather *et al.* (1991) that determined the efficacy of irradiation as a treatment for this pest in *Mangifera indica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J. 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L. 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J. 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R. 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Heather, N.W., Corcoran, R.J. & Banos, C. 1991. Disinfestation of mangoes with gamma irradiation against two Australian fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 84: 1304–1307.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M. 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M. 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L. 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A. 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

**Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for Bactrocea jarvisi* (2006-118)

2006-12 TPPT developed draft text

2007-05 SC approved draft text for MC

2007-10 Sent for MC under fast-track process

2008-07 TPPT revised draft text

2008-12 SC revised draft text for adoption via e-decision

2009-03 CPM-4 adopted Annex 4 to ISPM 28:2007

**ISPM 28. 2007: Annex 4 *Irradiation treatment for Bactrocera jarvisi* (2009).**  
Rome, IPPC, FAO.

Publication notes: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 5: Irradiation treatment for *Bactrocera tryoni* (2009)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 100 Gy minimum absorbed dose to prevent the emergence of adults of *Bactrocera tryoni* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Bactrocera tryoni</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Bactrocera tryoni</i> (Froggatt) (Diptera: Tephritidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Bactrocera tryoni</i> .

##### **Treatment schedule**

Minimum absorbed dose of 100 Gy to prevent the emergence of adults of *Bactrocera tryoni*.

Efficacy and confidence level of the treatment is ED<sub>99,9978</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18 (*Guidelines for the use of irradiation as a phytosanitary measure*).

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Bactrocera tryoni* (larvae and/or pupae) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Heather *et al.* (1991) that determined the efficacy of irradiation as a treatment for this pest in *Mangifera indica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Heather, N.W., Corcoran, R.J. & Banos, C.** 1991. Disinfestation of mangoes with gamma irradiation against two Australian fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 84: 1304–1307.
- Jessup, A.J., Rigney, C. J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

**Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for Bactrocera tryoni* (2006-119)

2006-12 TPPT developed draft text

2007-05 SC approved draft text for MC

2007-10 Sent for MC under fast-track process

2008-07 TPPT revised draft text

2008-12 SC revised draft text for adoption via e-decision

2009-03 CPM-4 adopted Annex 5 to ISPM 28:2007

**ISPM 28. 2007: Annex 5** *Irradiation treatment for Bactrocera tryoni* (2009). Rome, IPPC, FAO.

Publication notes: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 6: Irradiation treatment for *Cydia pomonella* (2009)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 200 Gy minimum absorbed dose to prevent the emergence of adults of *Cydia pomonella* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Cydia pomonella</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Cydia pomonella</i> (L.) (Lepidoptera: Tortricidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Cydia pomonella</i> .

##### **Treatment schedule**

Minimum absorbed dose of 200 Gy to prevent the emergence of adults of *Cydia pomonella*.

Efficacy and confidence level of the treatment is ED<sub>99,9978</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18 (*Guidelines for the use of irradiation as a phytosanitary measure*).

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Cydia pomonella* (larvae and/or pupae) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Mansour (2003) that determined the efficacy of irradiation as a treatment for this pest in *Malus domestica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

This phytosanitary treatment was adopted by the Fourth Session of the Commission on Phytosanitary Measures in 2009.

The annex is a prescriptive part of ISPM 28:2007.

---

#### **Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for Cydia pomonella* (2006-123)

2006-12 TPPT developed draft text

2007-05 SC approved draft text for MC

2007-10 Sent for MC under fast-track process

2008-07 TPPT revised draft text

2008-12 SC revised draft text for adoption e-decision

2009-03 CPM-4 adopted Annex 6 to ISPM 28:2007

**ISPM 28. 2007: Annex 6** *Irradiation treatment for Cydia pomonella* (2009). Rome, IPPC, FAO.

Publication notes: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 7: Irradiation treatment for fruit flies of the family Tephritidae (generic) (2009)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 150 Gy minimum absorbed dose to prevent the emergence of adults of fruit flies at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for fruit flies of the family Tephritidae (generic)
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	Fruit flies of the family Tephritidae (Diptera: Tephritidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of fruit flies of the family Tephritidae.

##### **Treatment schedule**

Minimum absorbed dose of 150 Gy to prevent the emergence of adults of fruit flies.

Efficacy and confidence level of the treatment is ED<sub>99,9968</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18 (*Guidelines for the use of irradiation as a phytosanitary measure*).

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable larvae and/or pupae during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Bustos *et al.* (2004), Follett & Armstrong (2004), Gould & von Windeguth (1991), Hallman (2004), Hallman & Martinez (2001), Hallman & Thomas (1999), Hallman & Worley (1999), Heather *et al.* (1991), Jessup *et al.* (1992), von Windeguth (1986) and von Windeguth & Ismail (1987) that determined the efficacy of irradiation as a treatment for this pest in *Averrhoa carambola*, *Carica papaya*, *Citrus paradisi*, *Citrus reticulata*, *Citrus sinensis*, *Lycopersicon esculentum*, *Malus domestica*, *Mangifera indica*, *Persea americana*, *Prunus avium* and *Vaccinium corymbosum*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J. 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Follett, P.A. & Armstrong, J.W. 2004. Revised irradiation doses to control melon fly, Mediterranean fruit fly, and Oriental fruit fly (Diptera: Tephritidae) and a generic dose for tephritid fruit flies. *Journal of Economic Entomology*, 97: 1254–1262.
- Gould, W.P. & von Windeguth, D.L. 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J. 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. 2004. Irradiation disinfestation of apple maggot (Diptera: Tephritidae) in hypoxic and low-temperature storage. *Journal of Economic Entomology*, 97: 1245–1248.
- Hallman, G.J. & Martinez, L.R. 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Hallman, G.J. & Thomas, D.B. 1999. Gamma irradiation quarantine treatment against blueberry maggot and apple maggot (Diptera: Tephritidae). *Journal of Economic Entomology*, 92: 1373–1376.
- Hallman, G.J. & Worley, J.W. 1999. Gamma radiation doses to prevent adult emergence from immatures of Mexican and West Indian fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 92: 967–973.

- Heather, N.W., Corcoran, R.J. & Banos, C.** 1991. Disinfestation of mangoes with gamma irradiation against two Australian fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 84: 1304–1307.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

#### Publication history

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for fruit flies of the family Tephritidae* (2006-126)

2006-12 TPPT developed draft text

2007-05 SC approved draft text for MC

2007-10 Sent for MC under fast-track process

2008-07 TPPT revised draft text

2008-12 SC revised draft text for adoption via e-decision

2009-03 CPM-4 adopted Annex 7 to ISPM 28:2007

**ISPM 28.** 2007: **Annex 7** *Irradiation treatment for fruit flies of the family Tephritidae* (2009). Rome, IPPC, FAO.

Publication notes: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 8: Irradiation treatment for *Rhagoletis pomonella* (2009)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 60 Gy minimum absorbed dose to prevent the development of phanerocephalic pupae of *Rhagoletis pomonella* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Rhagoletis pomonella</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Rhagoletis pomonella</i> (Walsh) (Diptera: Tephritidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Rhagoletis pomonella</i> .

##### **Treatment schedule**

Minimum absorbed dose of 60 Gy to prevent the development of phanerocephalic pupae of *Rhagoletis pomonella*.

Efficacy and confidence level of the treatment is ED<sub>99,9921</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18 (*Guidelines for the use of irradiation as a phytosanitary measure*).

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Rhagoletis pomonella* (larvae and/or pupae) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Hallman (2004) and Hallman & Thomas (1999) that determined the efficacy of irradiation as a treatment for this pest in *Malus domestica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J.** 2004. Irradiation disinfestation of apple maggot (Diptera: Tephritidae) in hypoxic and low-temperature storage. *Journal of Economic Entomology*, 97: 1245–1248.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Hallman, G.J. & Thomas, D.B.** 1999. Gamma irradiation quarantine treatment against blueberry maggot and apple maggot (Diptera: Tephritidae). *Journal of Economic Entomology*, 92: 1373–1376.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

**Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for pomonella* (2006-129)

2006-12 TPPT developed draft text

2007-05 SC approved draft text for MC

2007-10 Sent for MC under fast-track process

2008-07 TPPT revised draft text

2008-12 SC revised draft text for adoption via e-decision

2009-03 CPM-4 adopted Annex 8 to ISPM 28:2007

**ISPM 28. 2007: Annex 8** *Irradiation treatment for Rhagoletis pomonella* (2009).  
Rome, IPPC, FAO.

Publication notes: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 9: Irradiation treatment for *Conotrachelus nenuphar* (2010)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 92 Gy minimum absorbed dose to prevent the reproduction in adults of *Conotrachelus nenuphar* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Conotrachelus nenuphar</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Conotrachelus nenuphar</i> (Herbst) (Coleoptera: Curculionidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Conotrachelus nenuphar</i> .

##### **Treatment schedule**

Minimum absorbed dose of 92 Gy to prevent the reproduction in adults of *Conotrachelus nenuphar*.

Efficacy and confidence level of the treatment is ED<sub>99,9880</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18:2003.

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Conotrachelus nenuphar* (larvae, pupae and/or adults) during the inspection process. This does not imply a failure of the treatment.

Although the treatment may result in the presence of irradiated adults, the following factors may affect the likelihood of adults being found in traps in importing countries:

- Adults are rarely (if ever) present in shipped fruit because the insect pupates off the fruit.
- Irradiated adults are very unlikely to survive for more than one week, post-irradiation, and they are therefore less likely to spread than non-irradiated adults.

The Technical Panel on Phytopsanitary Treatments based its evaluation of this treatment on the research work undertaken by Hallman (2003) that determined the efficacy of irradiation as a treatment for this pest in *Malus domestica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2003. Ionizing irradiation quarantine treatment against plum curculio (Coleoptera: Curculionidae). *Journal of Economic Entomology*, 96: 1399–1404.
- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.

**von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

#### **Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for Conotrachelus nenuphar* (2006-120)

2006-12 TPPT developed draft text

2007-05 SC approved draft text for MC

2007-10 Sent for MC under fast-track process

2007-12 TPPT reviewed draft text

2008-12 SC revised draft text for adoption via e-decision

2009-03 Secretariat received formal objections prior to CPM-4

2009-05 SC requested to TPPT review

2009-11 TPPT reviewed and revised draft via email

2009-11 SC revised draft text for adoption

2010-03 CPM-5 adopted Annex 9 to ISPM 28:2007

**ISPM 28. 2007: Annex 9** *Irradiation treatment for Conotrachelus nenuphar* (2010). Rome, IPPC, FAO.

Publication history: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 10: Irradiation treatment for *Grapholita molesta* (2010)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 232 Gy minimum absorbed dose to prevent the emergence of adults of *Grapholita molesta* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Grapholita molesta</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Grapholita molesta</i> (Busck) (Lepidoptera: Tortricidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Grapholita molesta</i> .

##### **Treatment schedule**

Minimum absorbed dose of 232 Gy to prevent the emergence of adults of *Grapholita molesta*.

Efficacy and confidence level of the treatment is ED<sub>99,9949</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18:2003.

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Grapholita molesta* (larvae and/or pupae) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Hallman (2004) that determined the efficacy of irradiation as a treatment for this pest in *Malus domestica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

**Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added work topic *Irradiation treatment for Grapholita molesta* (2006-127A)

2006-12 TPPT developed draft text and recommended it to the SC

2007-07 SC revised draft text and approved for member consultation via email

2007-10 Member consultation under fast-track process

2008-07 TPPT reviewed and revised draft text via email

2008-12 SC revised draft text via e-decision

2009-03 Secretariat received formal objections prior to CPM-4

2009-05 SC requested TPPT for review draft text

2009-11 TPPT revised draft text via email

2009-11 SC reviewed draft text for adoption

2010-03 CPM-5 adopted Annex 10 to ISPM 28

**ISPM 28. 2007 Annex 10** *Irradiation treatment for Grapholita molesta* (2010).  
Rome, IPPC, FAO.

Publication history: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28 PHYTOSANITARY TREATMENTS**

#### **PT 11: Irradiation treatment for *Grapholita molesta* under hypoxia (2010)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 232 Gy minimum absorbed dose under hypoxic conditions to prevent oviposition of *Grapholita molesta* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Grapholita molesta</i> under hypoxia
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Grapholita molesta</i> (Busck) (Lepidoptera: Tortricidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Grapholita molesta</i> .

##### **Treatment schedule**

Minimum absorbed dose of 232 Gy to prevent oviposition of *Grapholita molesta*.

Efficacy and confidence level of the treatment is ED<sub>99,9932</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18:2003.

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Grapholita molesta* (larvae, pupae and/or adults) during the inspection process. This does not imply a failure of the treatment.

Although the treatment may result in the presence of irradiated adults, the following factors may affect the likelihood of adults being found in traps in importing countries:

- Only a very small percentage of adults are likely to emerge after irradiation.
- Irradiated adults are very unlikely to survive for more than one week, post-irradiation, and they are therefore less likely to spread than non-irradiated adults.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Hallman (2004) that determined the efficacy of irradiation as a treatment for this pest in *Malus domestica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

**Publication history**

*This is not an official part of the standard*

2006-04 CPM-1 added topic *Irradiation treatment for Grapholita melosa under hypoxia*

2006-12 TPPT developed draft text and recommended to the SC (2006-127B)

2007-07 SC revised draft text and approved for member consultation via email

2007-10 Member consultation under fast-track process

2008-07 TPPT reviewed and revised draft text via email

2008-12 SC revised draft text via e-decision

2009-03 Secretariat received formal objections prior to CPM-4

2009-05 SC requested the TPPT to review

2009-11 TPPT revised draft text via email

2009-11 SC revised draft text for adoption

2010-03 CPM-5 adopted Annex 11 to ISPM 28:2007

**ISPM 28. 2007: Annex 11** *Irradiation treatment for Grapholita molesta under hypoxia* (2010). Rome, IPPC, FAO.

Publication history: Last modified August 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28:2007 PHYTOSANITARY TREATMENTS**

#### **PT 12: Irradiation treatment for *Cylas formicarius elegantulus* (2011)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 165 Gy minimum absorbed dose to prevent the development of F1 adults of *Cylas formicarius elegantulus* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003 (*Guidelines for the use of irradiation as a phytosanitary measure*)<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment:</b>	Irradiation treatment for <i>Cylas formicarius elegantulus</i>
<b>Active ingredient:</b>	N/A
<b>Treatment type:</b>	Irradiation
<b>Target pest:</b>	<i>Cylas formicarius elegantulus</i> (Summers) (Coleoptera: Brentidae)
<b>Target regulated articles:</b>	All fruits and vegetables that are hosts of <i>Cylas formicarius elegantulus</i> .

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

### Treatment schedule

Minimum absorbed dose of 165 Gy to prevent the development of F1 adults of *Cylas formicarius elegantulus*.

Efficacy and confidence level of the treatment is ED99.9952 at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18:2003 (Guidelines for the use of irradiation as a phytosanitary measure).

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Cylas formicarius elegantulus* (eggs, larvae, pupae and/or adults) during the inspection process. This does not imply a failure of the treatment.

Countries with established trapping and surveillance activities for *Cylas formicarius elegantulus* need to take account of the fact that adult insects may be detected in the traps in the importing country. Although these insects will not establish, countries need to assess whether such treatments are applicable in their countries, i.e. whether or not such findings would disrupt existing surveillance programmes.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Follet (2006) and Hallman (2001) that determined the efficacy of irradiation as a treatment for this pest in *Ipomoea batatas*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos et al., 2004; Gould & von Windeguth, 1991; Hallman, 2004; Hallman & Martinez, 2001; Jessup et al., 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognised, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Follett, P.A.** 2006. Irradiation as a methyl bromide alternative for postharvest control of *Omphisa anastomosalis* (Lepidoptera: Pyralidae) and *Euscepes postfasciatus* and *Cylas formicarius elegantulus* (Coleoptera: Curculionidae) in sweet potatoes. *Journal of Economic Entomology*, 99: 32–37.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2001. Ionizing irradiation quarantine treatment against sweet potato weevil (Coleoptera: Curculionidae). *Florida Entomologist*, 84: 415–417.

- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

#### Publication history

*This is not an official part of the standard*

2006-12 TPPT developed draft text

2007-04 CPM-2 added topic *Irradiation treatment for Cylas formicarius elegantulus* (2006-124)

2007-10 SC revised draft text and approved for MC

2007-10 SC sent for MC under fast-track process

2008-03 Secretariat received formal objections prior to CPM-3

2008-08 SC revised draft text with TPPT consultation via email

2008-12 SC recommended draft text to CPM via e-decision

2009-03 Secretariat received formal objections prior to CPM-4

2009-05 SC requested TPPT to review

2009-08 TPPT revised draft text

2009-12 SC recommended draft text to CPM via e-decision

2010-03 Secretariat received formal objections prior to CPM-5

2010-05 SC requested TPPT to review

2010-07 TPPT revised draft text

2010-08 SC recommended draft text to CPM via e-decision

2011-03 CPM-6 adopted Annex 12 to ISPM 28

**ISPM 28.** 2007: **Annex 12** *Irradiation treatment for Cylas formicarius elegantulus* (2011). Rome, IPPC, FAO.

Publication history: Last modified December 2011



## INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES

### ISPM 28:2007 PHYTOSANITARY TREATMENTS

#### PT 13: Irradiation treatment for *Euscepes postfasciatus* (2011)

##### Scope of the treatment

This treatment applies to the irradiation of fruits and vegetables at 150 Gy minimum absorbed dose to prevent the development of F1 adults of *Euscepes postfasciatus* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003 (*Guidelines for the use of irradiation as a phytosanitary measure*)<sup>1</sup>.

##### Treatment description

<b>Name of treatment</b>	Irradiation treatment for <i>Euscepes postfasciatus</i>
<b>Active ingredient</b>	N/A
<b>Treatment type</b>	Irradiation
<b>Target pest</b>	<i>Euscepes postfasciatus</i> (Fairmaire) (Coleoptera: Curculionidae)
<b>Target regulated articles</b>	All fruits and vegetables that are hosts of <i>Euscepes postfasciatus</i> .

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

### Treatment schedule

Minimum absorbed dose of 150 Gy to prevent the development of F1 adults of *Euscepes postfasciatus*.

Efficacy and confidence level of the treatment is ED<sub>99,9950</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18:2003 (*Guidelines for the use of irradiation as a phytosanitary measure*).

This irradiation treatment should not be applied to fruit and vegetables stored in modified atmospheres.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live, but non-viable *Euscepes postfasciatus* (eggs, larvae, pupae and/or adults) during the inspection process. This does not imply a failure of the treatment.

Countries with established trapping and surveillance activities for *Euscepes postfasciatus* need to take account of the fact that adult insects may be detected in the traps in the importing country. Although these insects will not establish, countries need to assess whether such treatments are applicable in their countries, i.e. whether or not such findings would disrupt existing surveillance programmes.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Follett (2006) that determined the efficacy of irradiation as a treatment for this pest in *Ipomoea batatas*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica* and artificial diet) and *Grapholita molesta* (*Malus domestica* and artificial diet) (Bustos *et al.*, 2004; Gould & von Windeguth, 1991; Hallman, 2004, Hallman & Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth & Ismail, 1987). It is recognised, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Follett, P.A.** 2006. Irradiation as a methyl bromide alternative for postharvest control of *Omphisa anastomosalis* (Lepidoptera: Pyralidae) and *Euscepes postfasciatus* and *Cylas formicarius elegantulus* (Coleoptera: Curculionidae) in sweet potatoes. *Journal of Economic Entomology*, 99: 32–37.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.

- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

#### Publication history

*This is not an official part of the standard*

2006-12 TPPT developed draft text

2007-04 CPM-2 added the topic *Irradiation treatment for Euscepes postfasciatus* (2006-125)

2007-10 SC revised draft text and approved for MC

2007-10 SC sent for MC under fast-track process

2008-03 Secretariat received formal objections prior to CPM-3

2008-08 SC revised draft text with TPPT consultation via email

2008-12 SC recommended draft text to CPM via e-decision

2009-03 Secretariat received formal objections prior to CPM-4

2009-05 SC requested the TPPT to review

2009-08 TPPT revised draft text

2009-12 SC recommended draft text to CPM via e-decision

2010-03 Secretariat received formal objections prior to CPM-5

2010-05 SC requested TPPT to review

2010-07 TPPT revised draft text

2010-08 SC recommended draft text to CPM via e-decision

2011-03 CPM-6 adopted Annex 12 to ISPM 28

**ISPM 28.** 2007: **Annex 13** *Irradiation treatment for Euscepes postfasciatus* (2011). Rome, IPPC, FAO.

Publication history: Last modified December 2011



## **INTERNATIONAL STANDARDS FOR PHYTOSANITARY MEASURES**

### **ISPM 28:2007 PHYTOSANITARY TREATMENTS**

#### **PT 14: Irradiation treatment for *Ceratitis capitata* (2011)**

##### **Scope of the treatment**

This treatment applies to the irradiation of fruits and vegetables at 100 Gy minimum absorbed dose to prevent the emergence of adults of *Ceratitis capitata* at the stated efficacy. This treatment should be applied in accordance with the requirements outlined in ISPM 18:2003<sup>1</sup>.

##### **Treatment description**

<b>Name of treatment</b>	Irradiation treatment for <i>Ceratitis capitata</i>
<b>Active ingredient</b>	N/A
<b>Treatment type</b>	Irradiation
<b>Target pest</b>	<i>Ceratitis capitata</i> (Diptera: Tephritidae) (Mediterranean fruit fly)
<b>Target regulated articles</b>	All fruits and vegetables that are hosts of <i>Ceratitis capitata</i>

##### **Treatment schedule**

Minimum absorbed dose of 100 Gy to prevent the emergence of adults of *Ceratitis capitata*

Efficacy and confidence level of the treatment is ED<sub>99,9970</sub> at the 95% confidence level.

Treatment should be applied in accordance with the requirements of ISPM 18:2003.

---

<sup>1</sup> The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for approval of treatments. Treatments also do not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures prior to approval of a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

This irradiation treatment should not be applied to fruits and vegetables stored in modified atmospheres.

### Other relevant information

Since irradiation may not result in outright mortality, inspectors may encounter live but non-viable *Ceratitis capitata* (larvae and/or pupae) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research work undertaken by Follett and Armstrong (2004) and Torres-Rivera and Hallman (2007), which determined the efficacy of irradiation as a treatment for this pest in *Carica papaya* and *Mangifera indica*.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests (with hosts in parentheses): *Anastrepha ludens* (*Citrus paradisi* and *Mangifera indica*), *A. suspensa* (*Averrhoa carambola*, *Citrus paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*Citrus sinensis*, *Lycopersicon lycopersicum*, *Malus domestica*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus domestica*; also artificial diet) and *Grapholita molesta* (*Malus domestica*; also artificial diet) (Bustos *et al.*, 2004; Gould and von Windeguth, 1991; Hallman, 2004, Hallman and Martinez, 2001; Jessup *et al.*, 1992; Mansour, 2003; von Windeguth, 1986; von Windeguth and Ismail, 1987). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, then the treatment will be reviewed.

### References

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Follett, P.A. & Armstrong, J.W.** 2004. Revised irradiation doses to control melon fly, Mediterranean fruit fly, and Oriental fruit fly (Diptera: Tephritidae) and a generic dose for tephritid fruit flies. *Journal of Economic Entomology*, 97: 1254–1262.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2004. Ionizing irradiation quarantine treatment against Oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatments against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- ISPM 18.** 2003. Guidelines for the use of irradiation as a phytosanitary measure. Rome, IPPC, FAO.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. *Proceedings of the Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities*, 1990: 13–42.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.
- Torres-Rivera, Z. & Hallman, G.J.** 2007. Low-dose irradiation phytosanitary treatment against Mediterranean fruit fly (Diptera: Tephritidae). *Florida Entomologist*, 90: 343–346.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangoes. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.

**von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.

#### **Publication history**

*This is not an official part of the standard*

2007-12 TPPT developed draft text

2008-04 CPM-3 added topic *Irradiation treatment for Ceratitis capitata* (2007-204)

2008-11 SC revised draft text and approved for MC

2010-06 SC sent for MC under fast-track process

2010-12 SC recommended draft text to CPM via e-decision

2011-03 CPM-6 adopted Annex 14 to ISPM 28

**ISPM 28.** 2007: **Annex 14** *Irradiation treatment for Ceratitis capitata* (2011). Rome, IPPC, FAO.

Publication history: Last modified August 2011